

# Chemically optimized stereopure oligonucleotides direct ADAR-mediated RNA editing

Paloma H Giangrande, PhD  
Vice President, Platform Discovery Sciences  
Biology


*Oligonucleotides & Precision Therapeutics*  
March 15-16, 2022

WAVE™  
LIFE SCIENCES


# Forward looking statements

This document contains forward-looking statements. All statements other than statements of historical facts contained in this document, including statements regarding possible or assumed future results of operations, preclinical and clinical studies, business strategies, research and development plans, collaborations and partnerships, regulatory activities and timing thereof, competitive position, potential growth opportunities, use of proceeds and the effects of competition are forward-looking statements. These statements involve known and unknown risks, uncertainties and other important factors that may cause the actual results, performance or achievements of Wave Life Sciences Ltd. (the "Company") to be materially different from any future results, performance or achievements expressed or implied by the forward-looking statements. In some cases, you can identify forward-looking statements by terms such as "may," "will," "should," "expect," "plan," "aim," "anticipate," "could," "intend," "target," "project," "contemplate," "believe," "estimate," "predict," "potential" or "continue" or the negative of these terms or other similar expressions. The forward-looking statements in this presentation are only predictions. The Company has based these forward-looking statements largely on its current expectations and projections about future events and financial trends that it believes may affect the Company's business, financial condition and results of operations. These forward-looking statements speak only as of the date of this presentation and are subject to a number of risks, uncertainties and assumptions, including those listed under Risk Factors in the Company's Form 10-K and other filings with the SEC, some of which cannot be predicted or quantified and some of which are beyond the Company's control. The events and circumstances reflected in the Company's forward-looking statements may not be achieved or occur, and actual results could differ materially from those projected in the forward-looking statements. Moreover, the Company operates in a dynamic industry and economy. New risk factors and uncertainties may emerge from time to time, and it is not possible for management to predict all risk factors and uncertainties that the Company may face. Except as required by applicable law, the Company does not plan to publicly update or revise any forward-looking statements contained herein, whether as a result of any new information, future events, changed circumstances or otherwise.

# *SERPINA1* Z mutation: The most common cause of Alpha-1 antitrypsin deficiency (AATD)




# SERPINA1 Z mutation: The most common cause of Alpha-1 antitrypsin deficiency (AATD)




*Gain-of-function and loss-of-function disease*

## Liver Disease

Fibrosis → Cirrhosis → Hepatocellular Carcinoma



E342K mutation causes AAT proteotoxic stress, leading to progressive liver disease



## Lung Disease

Emphysema

Bronchiectasis




Low serum AAT leads to lung disease

# RNA base editing is uniquely suited to address the therapeutic goals for AATD


## Wave ADAR editing approach addresses goals of treatment:

**1) Restore** circulating, functional wild-type M-AAT




M-AAT reaches lungs to protect from proteases

**2) Reduce** Z-AAT protein aggregation in liver



Wild-type M-AAT protein replaces Z-AAT with RNA correction

**3) Retain** M-AAT physiological regulation




M-AAT secretion into bloodstream

## Alternative approaches address only a subset of treatment goals:

**Standard of care: protein augmentation** (11 $\mu$ M) addresses only lung manifestations

**siRNA approaches** address only liver disease


**Small molecule approaches** may address the lung and liver but do not generate wild-type M-AAT



~200K people in US and EU with mutation in *SERPINA1* Z allele (PI\*ZZ)

# Unlocking RNA editing with PRISM™ to develop GalNAc-AIMers: A-to-I editing oligonucleotides

Optimize AIMer design for endogenous transcripts and GalNAc conjugation




### ADAR enzymes

- First publication (1995) using oligonucleotide to edit RNA with endogenous ADAR<sup>1</sup>
- Catalyze conversion of A-to-I (G) in double-stranded RNA substrates
- ADAR1 is ubiquitously expressed across tissues, including liver and CNS
- Cellular reservoir of ADAR capacity supports directed editing in addition to homeostatic function

### Optimize AIMer design

- ✓ Substrate learnings from biology and structures
- ✓ Applied to oligonucleotides
- ✓ Applied PRISM chemistry

### PRISM-driven gains






← Improved potency

- PS/PO/PN
- PS/PO (Stereopure)
- PS/PO (Stereorandom)

# AIMers: Realizing potential of therapeutic RNA base editing by harnessing endogenous ADAR


Solved for key therapeutic attributes for potential best-in-class RNA editing therapeutics




- Systematized AIMer design enables rapid advancement of new targets
- Strong and broad IP in chemical and backbone modifications, stereochemistry patterns, novel and proprietary nucleosides

# Proof-of-concept RNA editing in NHP liver is durable and specific


**AIMers detected in liver of NHP at day 50**



**Substantial and durable editing in NHP liver *in vivo***



**ADAR editing with ACTB AIMer is highly specific**





# Proof-of-concept preclinical RNA editing data published in *Nature Biotechnology*

nature  
biotechnology

ARTICLES

<https://doi.org/10.1038/s41587-022-01225-1>

Check for updates

## Endogenous ADAR-mediated RNA editing in non-human primates using stereopure chemically modified oligonucleotides

Prashant Monian<sup>1,2</sup>, Chikdu Shivalila<sup>1,2</sup>, Genliang Lu<sup>1</sup>, Mamoru Shimizu<sup>1</sup>, David Boulay<sup>1</sup>, Karley Bussow<sup>1</sup>, Michael Byrne<sup>1</sup>, Adam Bezigian<sup>1</sup>, Arindom Chatterjee<sup>1</sup>, David Chew<sup>1</sup>, Jigar Desai<sup>1</sup>, Frank Favalaro<sup>1</sup>, Jack Godfrey<sup>1</sup>, Andrew Hoss<sup>1</sup>, Naoki Iwamoto<sup>1</sup>, Tomomi Kawamoto<sup>1</sup>, Jayakanthan Kumarasamy<sup>1</sup>, Anthony Lamattina<sup>1</sup>, Amber Lindsey<sup>1</sup>, Fangjun Liu<sup>1</sup>, Richard Looby<sup>1</sup>, Subramanian Marappan<sup>1</sup>, Jake Metterville<sup>1</sup>, Ronelle Murphy<sup>1</sup>, Jeff Rossi<sup>1</sup>, Tom Pu<sup>1</sup>, **Bijay Bhattarai**<sup>1</sup>, Stephany Standley<sup>1</sup>, Snehlata Tripathi<sup>1</sup>, Hailin Yang<sup>1</sup>, Yuan Yin<sup>1</sup>, Hui Yu<sup>1</sup>, **Cong Zhou**<sup>1</sup>, Luciano H. Apponi<sup>1</sup>, Pachamuthu Kandasamy<sup>1</sup> and **Chandra Vargeese**<sup>1</sup>✉

Technologies that recruit and direct the activity of endogenous RNA-editing enzymes to specific cellular RNAs have therapeutic potential, but translating them from cell culture into animal models has been challenging. Here we describe short, chemically modified oligonucleotides called AIMers that direct efficient and specific A-to-I editing of endogenous transcripts by endogenous adenosine deaminases acting on RNA (ADAR) enzymes, including the ubiquitously and constitutively expressed ADAR1 p110 isoform. We show that AIMers achieve up to 50% editing with no bystander editing of the endogenous AC1B transcript in non-human primates.

**Specificity *in vitro* & *in vivo* (NHPs)**


***In vitro-in vivo* translation (NHPs)**

**GaINac conjugation**


**Foundational AIMer SAR**

# Robust SERPINA1 RNA base editing in mouse model for AATD restores M-AAT protein in serum

**RNA editing in mice  
day 7 (3x5 mg/kg, SC)**




**AAT serum protein concentration in mice  
day 7 (3x10 mg/kg, SC)**



huADAR/SERPINA1 mice


# Bi-weekly AIMer treatment results in serum AAT protein levels in mice above anticipated therapeutic threshold

GalNAc-AIMer results in serum AAT protein levels >11  $\mu\text{M}$  at week 19 in transgenic mice




# Bi-weekly AIMer treatment supports robust RNA editing and M-AAT protein expression in mice

GalNAc-AIMER results in serum AAT protein levels >11 uM at week 19 in transgenic mice




~60% RNA editing & ~70% serum M-AAT protein (week 19 data)



huADAR/SERPINA1 mice

# Further analyses suggest functional effects in mouse liver at 19 weeks



GalNAc-AIMER results in serum AAT protein levels >11  $\mu$ M at week 19 in transgenic mice



huADAR/SERPINA1 mice


↑ 10 mg/kg AIMER dose

~60% RNA editing & ~70% serum M-AAT protein (week 19 data)




# AIMer-directed editing is highly specific in mice; no bystander editing observed on SERPINA1 transcript

**RNA editing only detected at PiZ mutation site in SERPINA1 transcript (mouse liver)**




**RNA editing across transcriptome (mouse liver)**



# Opportunity for novel and innovative AIMer therapeutics


## Correct driver mutations with AIMers

Free-uptake of chemically modified oligonucleotides



**Endogenous ADAR**

## Restore or correct protein function



## Upregulate expression

## Modify function

## Modulate protein-protein interaction

## Post-translational modification

## Alter folding or processing

## Modulate protein interactions with AIMers

### **Examples**

AATD

Rett syndrome

Recessive or dominant  
genetically defined  
diseases

### **Examples**

Haploinsufficient  
diseases

Loss of function

Neuromuscular

Dementias

Familial epilepsies


Neuropathic pain

# Efficient and durable editing in mouse CNS with unconjugated AIMer



huADAR mice

Peak editing observed 4-weeks post-single ICV dose across tissues



	Cortex	Hippocampus	Striatum	Brain stem	Cerebellum	Spinal cord
Peak editing	30%	>40%	25%	>40%	50%	>65%

Transgenic huADAR mice were administered 100  $\mu$ g AIMer or PBS on day 0 and evaluated for UGP2 editing across CNS tissues at 1, 4, 8, 12 and 16-weeks post dose. Percentage UGP2 editing determined by Sanger sequencing. Stats: 2-way ANOVA with post-hoc comparison to PBS (n=5 per time point per treatment) \*P<0.05, \*\*P<0.01, \*\*\*P<0.001, \*\*\*\*P<0.0001. ICV intracerebroventricular; PBS phosphate buffered saline



# Acknowledgements



- *Colleagues and contributors from Wave Life Sciences*

## Endogenous ADAR-mediated RNA editing in non-human primates using stereopure chemically modified oligonucleotides

Prashant Monian<sup>1,2</sup>, Chikdu Shivalila<sup>1,2</sup>, Genliang Lu<sup>1</sup>, Mamoru Shimizu<sup>1</sup>, David Boulay<sup>1</sup>, Karley Bussow<sup>1</sup>, Michael Byrne<sup>1</sup>, Adam Bezigian<sup>1</sup>, Arindom Chatterjee<sup>1</sup>, David Chew<sup>1</sup>, Jigar Desai<sup>1</sup>, Frank Favalaro<sup>1</sup>, Jack Godfrey<sup>1</sup>, Andrew Hoss<sup>1</sup>, Naoki Iwamoto<sup>1</sup>, Tomomi Kawamoto<sup>1</sup>, Jayakanthan Kumarasamy<sup>1</sup>, Anthony Lamattina<sup>1</sup>, Amber Lindsey<sup>1</sup>, Fangjun Liu<sup>1</sup>, Richard Looby<sup>1</sup>, Subramanian Marappan<sup>1</sup>, Jake Metterville<sup>1</sup>, Ronelle Murphy<sup>1</sup>, Jeff Rossi<sup>1</sup>, Tom Pu<sup>1</sup>, Bijay Bhattarai<sup>1</sup>, Stephany Standley<sup>1</sup>, Snehlata Tripathi<sup>1</sup>, Hailin Yang<sup>1</sup>, Yuan Yin<sup>1</sup>, Hui Yu<sup>1</sup>, Cong Zhou<sup>1</sup>, Luciano H. Apponi<sup>1</sup>, Pachamuthu Kandasamy<sup>1</sup> and Chandra Vargese<sup>1,2</sup>✉

Technologies that recruit and direct the activity of endogenous RNA-editing enzymes to specific cellular RNAs have therapeutic potential, but translating them from cell culture into animal models has been challenging. Here we describe short, chemically modified oligonucleotides called AIMers that direct efficient and specific A-to-I editing of endogenous transcripts by endogenous adenosine deaminases acting on RNA (ADAR) enzymes, including the ubiquitously and constitutively expressed ADAR1 p110 isoform. We show that fully chemically modified AIMers with chimeric backbones containing stereopure phosphorothioate and nitrogen-containing linkages based on phosphoryl guanidine enhanced potency and editing efficiency 100-fold compared with those with uniformly phosphorothioate-modified backbones in vitro. In vivo, AIMers targeted to hepatocytes with N-acetylgalactosamine achieve up to 50% editing with no bystander editing of the endogenous ACTB transcript in non-human primate liver, with editing persisting for at least one month. These results support further investigation of the therapeutic potential of stereopure AIMers.

Recruiting endogenous RNA-editing enzymes using chemically modified oligonucleotides holds promise for treating human disease. The most common mutation in human genes is transition from cytosine (C) to thymine (T), and CpG dinucleotides are well established hot spots for disease-causing mutations. The ADAR family of enzymes catalyze adenine (A)-to-inosine (I) changes in the transcriptome<sup>1</sup>. Because I is read as guanine (G) by the translational machinery<sup>2</sup>, ADAR-mediated RNA editing has the potential to revert these disease-causing transitions at the RNA level. The potential scope for application of A-to-I editing is large, including modulation of polar or charged amino acids, stop codons or RNA regulatory sequences<sup>3,4</sup>, eliciting diverse functional outcomes (for example, restored protein expression or function). Chemical modifications are known to confer drug-like properties to oligonucleotides. We set out to determine whether control over backbone chemistry and stereochemistry and other chemical modifications to an oligonucleotide (Fig. 1 and Supplementary Note 1) can be optimized to elicit sequence-specific A-to-I RNA editing with endogenous ADAR enzymes. As therapeutics, reversible RNA editing with oligonucleotides may represent a safer option than those that edit genomic DNA. Early technologies designed to elicit RNA editing in vitro required an exogenous enzyme and an oligonucleotide<sup>5-17</sup>. These approaches led to overexpression of editing enzyme and substantial off-target editing<sup>6,18-19</sup>. Recent advances have overcome the need for exogenous enzymes in vitro<sup>20</sup>, but they still use long oligonucleotides that require ancillary delivery reporter and exogenous ADAR enzyme in the presence or absence of cell culture. So far, these technologies have yielded nominal editing in vivo<sup>21</sup>. Leveraging our oligonucleotide chemistry platform, we developed relatively short oligonucleotides that elicit A-to-I RNA editing with high efficiency using endogenous ADAR enzymes. These oligonucleotides, called AIMers, are short and fully chemically modified with stereopure phosphorothioate (PS) and nitrogen-containing linkages based on phosphoryl guanidine. In vitro, they enhanced potency and A-to-I editing efficiency compared to uniformly PS-modified AIMers, and in vivo, N-acetylgalactosamine (GalNAc)-modified AIMers achieved up to 50% editing with no bystander editing in non-human primate (NHP) liver that persisted for at least 1 month.

**AIMers support RNA editing.** To evaluate RNA-editing efficiency in mammalian cells, we created a luciferase reporter gene with endogenous Gluc and *Cypridina* (Cluc). In the absence of editing with oligonucleotides, only Gluc is expressed, whereas A-to-I editing permits expression of Cluc, providing a measure of RNA-editing efficiency and protein expression (Extended Data Fig. 1a). AIMers were designed to mimic naturally occurring double-stranded RNA ADAR substrates, as in the GluR2 transcript<sup>6,10,20</sup> (Extended Data Fig. 1b). To benchmark RNA editing, we transfected 293T cells with the reporter and exogenous ADAR enzyme in the presence or absence